

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This is the peer-reviewed, manuscript version of an article published in the *Journal of Equine Veterinary Science*. The version of record is available from the journal site:

<https://doi.org/10.1016/j.jevs.2020.103109>.

© 2019. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Acute life-threatening laryngeal dysfunction in a draft horse recovering from general anesthesia: a case report

AUTHORS: Hayley Linda Ronaldson, Paolo Monticelli, Roger Smith, Chiara Adami

JOURNAL: Journal of Equine Veterinary Science

PUBLISHER: Elsevier

PUBLICATION DATE: 25 April 2020

DOI: 10.1016/j.jevs.2020.103109

Acute life-threatening laryngeal dysfunction in a draft horse recovering from general anesthesia: a case report

Hayley Linda Ronaldson, Paolo Monticelli, Roger Smith and Chiara Adami

Department of Clinical Sciences and Services, Royal Veterinary College, University of London, Hatfield, AL9 7TA United Kingdom

Corresponding author

Paolo Monticelli

Department of Clinical Sciences and Services, Royal Veterinary College, University of London, Hatfield AL9 7TA United Kingdom

Email: pmonticelli@rvc.ac.uk

Phone: 0044 (0) 7476794355

Abstract

A 13-year-old Shire horse was anesthetized for an elective orthopaedic procedure. During recovery from anesthesia, the occurrence of severe acute dyspnea required a second anesthetic to allow endoscopy-guided nasotracheal intubation. Endoscopic findings were decreased mobility and swelling of the arytenoids with narrowing of the laryngeal *aditus*. Owing to a dislodgement of the nasotracheal tube during recovery, a third anesthetic was carried out to perform emergency tracheostomy. Recovery from the third anesthetic was long and the horse developed a post-anesthetic myopathy. The clinical conditions improved during the following 48 hours, and an endoscopic examination performed prior to discharge revealed unremarkable laryngeal function. It was hypothesized that mechanical stimulation of the trachea during the phases of intubation and extubation caused traumatic laryngeal dysfunction, and that draft horses might require additional care during the perioperative period.

Keywords: Anesthesia, draft horse, larynx, perioperative complication

1. Introduction

Equine anesthesia is associated with an incidence of fatalities of 1.1% [1]. Compared to other domestic mammals, horses have a higher risk for peri-anesthetic complications, of which bone fractures, myopathies and respiratory impairment have been frequently reported [1-3]. Among respiratory complications, upper airway obstruction and pulmonary edema have been described in equine patients [4-6]. The former mainly occurs after extubation, as a result of either nasal obstruction by secretions or edema, or displacement of the soft palate owing to a combination of mechanical and pharmacological effects [7]. Regarding pulmonary edema, this complication is most commonly the ultimate result of upper airway obstruction. In an attempt to overcome the non-patent airways, large negative intra-thoracic pressures are generated, resulting in movement of fluid from the vascular compartment to the interstitium and, ultimately, into the alveolar space [3, 8-11]. Other causes of peri-anesthetic pulmonary edema, including microembolism following orthopaedic surgery and side effects of opioids and alpha 2- adrenoreceptor agonists [12,13], have also been reported in horses.

This report describes the management of a life-threatening upper airways complication and post-anesthetic myopathy in a draft horse recovering from general anesthesia for an elective procedure. The occurrence of transient post-anesthetic laryngeal dysfunction in the absence of pre-existing respiratory or neurological conditions has been previously documented in horses [4,5]. Nevertheless, to the best of the authors' knowledge, cases that resolved within 24 hours have not been reported yet.

2. Case description

A 13-year-old female Shire was referred to the Equine Referral Hospital of the Royal Veterinary College for evaluation of an acute onset lameness of the right hind limb associated with digital sheath effusion of three weeks duration. On examination, lameness was mild and was resolved with intra-theal diagnostic analgesia of the digital flexor tendon sheath (DFTS). A concurrently performed contrast tenogram showed no obvious abnormality of the *manica flexoria* but did suggest the presence of a constricted plantar annular ligament because contrast did not pass freely into the proximal pouch of the DTFS after injection via the distal pouch. Ultrasonographic examination of the DFTS revealed marked synovial thickening and effusion of the tendon sheath but no other significant abnormalities. Tenoscopy under general anesthesia was scheduled for the next day upon request of the orthopaedic surgeon.

Pre-anesthetic physical examination revealed heart rate (HR) of 40 beats minute⁻¹, respiratory rate (fR) of 12 breaths minute⁻¹, rectal temperature of 37°C, and pink mucous membranes with capillary refill time less than 2 seconds. Body weight and body condition score were 694 kg and 3/5 [14], respectively. Chest auscultation and blood results were unremarkable. An American Society of Anaesthesiologists category of 1 was assigned to the horse; pre-operative withholding time for food was 12 hours whereas water was made available until premedication.

Intramuscular (IM) acepromazine (AceSedate, Jurox, UK; 0.02 mg kg⁻¹) was administered as a tranquilizer, 60 minutes before placement of a 14 G catheter in the left jugular vein. The horse was then transferred from the stable to the induction box, where it received intravenous (IV) romifidine (Sedivet, Boehringer Ingelheim, UK; 0.04 mg kg⁻¹) and morphine (Morphine Sulphate, Mercury Pharmaceuticals, Ireland; 0.1 mg kg⁻¹) as premedication. General anesthesia was induced with IV ketamine (Ketamidol, Chanelle, UK; 2.2 mg kg⁻¹) and midazolam (Dormicum, Roche, UK; 0.05 mg kg⁻¹); induction was smooth and uneventful. Additional ketamine (1 mg kg⁻¹) was administered prior to attempting endotracheal

intubation, owing to the presence of nystagmus and poor myorelaxation that were considered by the anesthesiologist in charge to be indicative of superficial anesthetic depth. A 24 mm internal diameter endotracheal (ET) tube was then inserted into the trachea via the oro-tracheal route and cuffed with a syringe to ensure air-tight seal. Following the patient was transported to theatre by an overhead crane system where it was positioned in dorsal recumbency on the surgical table with the head and neck slightly elevated to avoid their possible over-extension. The ET tube was connected to a circle breathing system, and isoflurane (IsoFlo, Abbott, UK) was delivered in a mixture of oxygen and medical air (1:1). Intermittent positive pressure ventilation was initiated at a rate of 8 breaths/minute with initial setting of 6.5 L and 17 cm H₂O as tidal volume (TV) and peak inspiratory pressure (PIP), respectively. The horse was instrumented with a multiparametric monitor that included electrocardiography, capnography, pulse oximetry and continuous measurement of invasive arterial blood pressure (via facial artery) and oesophageal temperature. Physiological variables were manually recorded on the anesthetic record every 5 minutes. Arterial blood was sampled from an indwelling 21G catheter, placed in the facial artery for continuous blood pressure measurement, and analyzed at 30-60-minute intervals with a portable device (i-STAT blood analyzer and cartridges CG4+; Abbott, NJ, USA). During surgery, the horse received IV fluids (Hartmann's solution, Fresenius Kabi, UK; 5 mL kg hr⁻¹) as well as a continuous rate of infusion (CRI) of romifidine (0.04 mg kg hr⁻¹). A CRI of dobutamine (Dobutamine Concentrate, Hameln Pharmaceutical, Germany; 0.5-1 mcg kg minute⁻¹) was initiated before successful arterial catheter placement owing to poor peripheral pulse quality, as judged subjectively by the anesthesiologist in charge.

Tenoscopic examination of the DFTS revealed the presence of a small tear on the lateral border of the *manica flexoria*, as well as constriction of the plantar annular ligament of the fetlock canal; these lesions were treated with resection of the *manica flexoria* and annular ligament desmotomy, respectively.

General anesthesia was without critical events. Transient hypotension (mean arterial pressure less than 70 mmHg) occurred during surgery; however, it was promptly detected and successfully treated with dobutamine infusion. Total anesthesia and surgery time were 200 and 105 min, respectively.

Recovery from anesthesia was assisted, with head-tail-rope method. At recovery, airflow was detected from both nares when the cuff of the ET tube was deflated and judged subjectively as adequate by the anesthesiologist in charge. The trachea of the horse was extubated after spontaneous swallowing resumed. The horse showed an unremarkable breathing pattern during left lateral recumbency, which lasted for approximately 30 minutes. Thereafter, after one unsuccessful attempt to stand, the horse managed to achieve sternal recumbency; at this point, the occurrence of sudden onset severe inspiratory dyspnoea, characterized by marked inspiratory stridor and visible cyanosis of the oral mucous membranes, required emergency reintubation. For this purpose, general anesthesia was induced with IV ketamine (2.5 mg kg⁻¹). Blind orotracheal intubation of the trachea with a 24 mm ET tube was unsuccessfully attempted; it was therefore decided to use endoscopic guidance. Endoscopy revealed swelling of the larynx resulting in narrowing of the laryngeal *aditus*, as well as absence of arytenoid movement during breathing, which suggested bilateral laryngeal dysfunction, possibly as a result of mechanical trauma. The soft palate was not displaced. Nasotracheal intubation was performed under endoscopic guidance, with an 18 mm ET tube, with the intention of leaving the ET tube in place after standing. However, during recovery from anesthesia, while the horse attempted to stand the nasotracheal tube was dislodged and kinked, and severe dyspnea reoccurred. This complication required immediate interventions; a third anesthetic was carried out to allow emergency tracheostomy in the recovery box. General anesthesia was induced with IV ketamine (2.5 mg kg⁻¹),

following xylazine (Chanazine, Channele, UK; 0.05 mg kg⁻¹, IV) premedication. Endoscopy-guided orotracheal intubation was accomplished with a 22 mm ET tube that was removed once the surgical procedure was completed and the tracheostomy tube was in place and secured. At this point, the horse was allowed to recover. Recovery from anesthesia was protracted and the horse remained in sternal recumbency for approximately two hours. In order to treat possible hypotension, suspected based on poor peripheral pulse, 2 L of hypertonic 7.2% NaCl and ephedrine (Ephedrine hydrochloride, Hameln Pharmaceutical, Germany; 0.05 mg kg⁻¹) were administered IV. At physical exam, rectal temperature was 35.5°C, pulse rate was 40 beats minute⁻¹ and respiratory rate was 10 breaths minute⁻¹. Hypothermia was treated with the use of blankets and infrared lamp heaters. Blood results from a sample collected during recovery revealed increased serum aspartate transaminase (676 U/L; reference interval: 198-476 U/L) and creatine kinase (15088 U/L; reference interval: 133-738 U/L). Based on these findings, a myopathy was suspected as further post-anesthetic complication. After three hours in the recovery box and five unsuccessful attempts to stand, an Anderson Sling was used to assist and maintain standing.

During the following 12 hours, the horse was closely monitored in the Intensive Care Unit of the hospital; support therapy consisted of IV fluids (Hartmann's solution, 2 mL kg hr⁻¹), analgesics (IV phenylbutazone, 4.4 mg kg⁻¹ SID, and IM morphine, 0.1 mg kg⁻¹ every four hours). Additionally, for the first two hours after standing oxygen was supplemented, via tracheostomy tube, at variable rate (2-4 L min⁻¹); thereafter, an arterial blood sample collected from the external carotid artery while the horse was breathing room air revealed normal blood gas results (arterial partial pressure of oxygen [PaO₂]: 98 mmHg, arterial oxygen saturation [SaO₂]: 97%; arterial partial pressure of carbon dioxide [PaCO₂]: 48 mmHg). Oxygen supplementation was discontinued; breathing remained unremarkable and the clinical condition of the horse was deemed good.

An endoscopic re-examination repeated 24 hours after the third anesthetic revealed normal abduction of the arytenoids. Thereafter, the tracheostomy tube was first blocked off to verify the laryngeal function, and then removed as dyspnea had not occurred. The surgical wound at the tracheostomy site was allowed to heal by second intention. Phenylbutazone administration was continued in the postoperative period for a total of five days.

The patient was discharged from the hospital 48 hours after recovery from the third anesthetic, and then reexamined after eight weeks. During this time, the horse owner did not notice any sign of respiratory dysfunction.

3. Discussion

This report describes the occurrence of acute, severe laryngeal dysfunction, confirmed by endoscopic findings, as a potentially life-threatening peri-anesthetic complication in a healthy horse undergoing anesthesia for an elective surgical procedure. The narrowing of the laryngeal *aditus* occurred without complete closure of the glottis; moreover, it was noticed that the arytenoids remained stationary in a neutral position throughout the respiratory cycle. These findings seem to suggest that, even if some degree of laryngospasm might have occurred, most likely the primary cause of the acute life-threatening dyspnea observed in this horse was rather laryngeal hypomotility or paralysis, a complication that has been described before in horses recovering from anesthesia [4,5]. Since the laryngeal dysfunction resolved spontaneously within 24 hours, and considering that the horse had not suffered from respiratory disease either prior to anesthesia, or during the eight weeks after that, it would be reasonable to assume that the condition was the result of a mechanical trauma or stimulation to the arytenoids, possibly occurring during tracheal intubation/extubation. In support of this hypothesis, the laryngeal swelling detected on endoscopic examination would suggest a

recent traumatic injury. It is however unclear how laryngeal trauma occurred in the horse of this report. Orotracheal intubation was performed by an Anesthesia resident in-training, supervised by a recognized specialist in Veterinary Anesthesia. Furthermore, a relatively small endotracheal tube was used, which should have decreased the risk of iatrogenic mechanical injury. Although orotracheal placement of the ET tube was without major complications, mild resistance was encountered at first attempt, which required reinsertion of the ET tube after repositioning of the head and neck. It is hypothesized that repeated, although gentle, manipulation of the airways was sufficient to cause laryngeal trauma that might have affected the normal laryngeal function of this horse.

Laryngospasm seemed unlikely also based on the onset of the dyspnea. Laryngospasm is a transient dysfunction of the arytenoids, usually triggered by peri-glottic stimulation that activates the vagus nerve and characterized by prolonged contraction of the intrinsic laryngeal muscles, which results in partial or complete airways closure [15-17]. This condition has been reported in horses and cats following mechanical stimulation associated with tracheal intubation [16-18], and in humans as a postoperative complication during removal of the ET tube [19,20]. Literature suggests that, in human patients, post-anesthetic laryngospasm is more likely to occur in the presence of underlying neurological or upper respiratory diseases [19,21]. Extubation-induced laryngospasm is likely to cause signs of respiratory impairment shortly after extubation, whereas in the horse of this report the onset of the dyspnea was approximately 30 minutes after extubation.

Underlying neurological or respiratory conditions were not suspected prior to anesthesia based on history and clinical findings; nevertheless, it should be emphasized that idiopathic laryngeal hemiplegia is overrepresented in draft horses, in which a prevalence of 35% has been reported [22]. Factors predisposing draft horses to laryngeal dysfunction may be the long neck and large body size, potentially causing a greater tension on the left recurrent laryngeal nerve, which in turn might compromise the blood supply to the larynx and result in nerve damage [23]. Additionally, working with a flexed head and neck carriage may possibly cause decreased degree of arytenoid abduction and atrophy of the muscles surrounding the laryngeal saccules [24-26]. Although unlikely, the presence of an underlying, subclinical laryngeal disease that may have predisposed the horse to peri-anesthetic respiratory complications cannot be completely excluded.

The increase in serum creatine kinase, together with the observed muscular weakness and inability of the horse to stand, suggests that post-anesthetic myopathy may have occurred as further complication. An association between the post-anesthetic myopathy (PAM) and acute laryngeal dysfunction, although unlikely also based on previously published literature, cannot be completely ruled out [5].

Post-anesthetic myopathies represent a life-threatening complication more likely to occur in heavy and large horses, and after prolonged anesthetics resulting in prolonged compression of the *logissimus dorsi* and gluteal muscles, and less commonly, of the *adductor* group when the horse is positioned in dorsal recumbency [27]. The horse of this report was relatively large, and the total recumbency time, during which muscular compression could have occurred, was more than six hours. Because the muscular weakness resolved spontaneously and the clinical condition of the horse continued to improve during the days following surgery, it is believed that the PAM was not generalized but affected some isolated muscle groups. Another condition that was not possible to exclude was the Equine Polysaccharide Storage Myopathy (EPSM), an inheritable disease characterized by alterations of glycogen storage and possibly over-represented in the draft horse [28]. It has been reported that EPSM can cause 'PAM-like' symptoms similar to the ones observed in the horse of this report [29]. Ideally, EPSM screening tests and a complete ultrasonographic examination could have been useful to make a more precise diagnosis through identification

of the affected muscle [30]; nevertheless, further diagnostics was declined by the horse owner owing to restricted budget.

Reduced anesthetic time and correct positioning on the surgical table are important factors to minimize the risk of morbidities associated with equine anesthesia [31]. Previous reports suggest that there might be an association between postoperative upper airways obstructions and either dorsal recumbency [4,6], or extension of the neck [5,6], or both, during anesthesia. Abrahamsen and colleagues [4] hypothesized that extension of the neck during anesthesia might alter the blood supply to the local anatomical structures, potentially leading to nerve damage. The horse of this report, although it had the head slightly elevated to avoid over-extension of the neck during anesthesia, was positioned in dorsal recumbency for several hours. The reasons for such long duration of anesthesia and surgery were the involvement of residents-in-training, which is common in teaching Institutions, and the fact that an emergency colic surgery had to be initiated while the horse of the report was already being anaesthetized – an inconvenience that decreased the availability of technicians and other non-medical personnel in the operating theatre. Whenever possible, optimization of time and resources is strongly advisable in order to decrease the anesthetic time and therefore the risk of anesthesia-related complications in equine patients.

Prompt recognition of the complication, rapid interventions and teamwork were essential for the good outcome of this case. One critical moment was the induction of the second anesthetic to allow emergency nasotracheal intubation, which required entrance into the recovery box while the horse was very excitable and struggled to breathe. The rationale for placing a nasotracheal tube at that time was because it was believed that, once conscious, the horse would have tolerated a nasotracheal tube better than an orotracheal one, which would have ensured maintenance of the airways' patency during recovery. However, the presence of the tube did not allow a proper endoscopic evaluation of the larynx prior to extubation, a diagnostic test which, with hindsight, would have been useful in this case. It is therefore sensible to state that performing an emergency tracheostomy immediately, once a patent airway had been established, would have been a safer approach to manage this case.

4. Conclusions

In conclusion, this report suggests that draft horses might require additional care during the perioperative period. The optimization of the anesthetic management and surgical time are essential in horses that for size and body weight may be prone to develop PAM or other anesthesia-related complications.

Additionally, ensuring prompt availability of a portable endoscope as well as of a surgical kit to perform emergency tracheostomy are essential for successful treatment of upper airway dysfunction during the peri-anesthetic period.

Acknowledgments

The authors thank all the clinicians involved in the management of the case.

References

- [1] Dugdale AHA, Obhrai J, Cripps PJ. Twenty years later: a single-centre, repeat retrospective analysis of equine perioperative mortality and investigation of recovery quality. *Vet Anaesth Analg* 2016;43:171-78.
- [2] Johnston GM, Eastment JK, Wood JLN, Taylor PM. The confidential enquiry into perioperative equine fatalities (CEPEF): mortality results of Phases 1 and 2. *Vet Anaesth Analg* 2002;29:159-70.
- [3] Senior M. Post-anaesthetic pulmonary oedema in horses: a review. *Vet Anaesth Analg* 2005;32:193-200.
- [4] Abrahamsen EJ, Bohanon TC, Bednarski RM, Hubbell JA, Muoir WW. Bilateral arytenoid cartilage paralysis after inhalation anesthesia in a horse. *J Am Vet Med Assoc* 1990;197:1363-1365.
- [5] Dixon PM, Railton DI, McGorum BC. Temporary bilateral laryngeal paralysis in a horse associated with general anaesthesia and post anaesthetic myositis. *Vet Rec* 1993;132:29-32.
- [6] Ball MA, Trim CM (1996) Post-anaesthetic pulmonary oedema in two horses. *Equine vet. Educ* 1996;8:13-16.
- [7] Wagner AE. Complications in equine anesthesia. *Vet Clin North Am Equine Pract* 2009;24:735-52.
- [8] Hackett RP, Ducharme NG, Ainsworth DM, Erickson BK, Erb HN, Soderholm LV, et al. Effects of extrathoracic airway obstruction on intrathoracic pressure and pulmonary artery pressure in exercising horses. *Am J Vet Res* 1999;60:485-94.
- [9] Holcombe SJ, Derksen FJ, Stick JA, Robinson NE, Boehler DA. Effect of nasal occlusion on tracheal and pharyngeal pressures in horses. *Am J Vet Res* 1996;57:1258-60.
- [10] Jackson JA, Ducharme NG, Hackett RP, et al. Effects of airway obstruction on transmural pulmonary artery pressure in exercising horses. *Am J Vet Res* 1997;58:897-903.
- [11] Tute AS, Wilkins PA, Gleed RD, Credille KM, Murphy DJ, Ducharme NG. Negative pressure pulmonary edema as a post-anesthetic complication associated with upper airway obstruction in a horse. *Vet Surg* 1996;25:519-23.
- [12] Jones RS, Payne-Johnson CE, Seymour CJ. Pulmonary micro-embolism following orthopaedic surgery in a Thoroughbred gelding. *Equine Vet J* 1988;20:382-84.
- [13] Shaw ML, Carpenter JW, Leith DE. Complications with the use of carfentanil citrate and xylazine hydrochloride to immobilize domestic horses. *J Am Vet Med Assoc* 1995;206:833-36.
- [14] Carroll CL, Huntington PJ. Body condition scoring and weight estimation of horses. *Equine Vet J* 1988;20:41-5.
- [15] Gavel G, Walker RWM. Laryngospasm in anaesthesia. *CEACCP* 2014;14:47-51.
- [16] Goulden BE, Barnes GR, Quinlan TJ. A case of equine laryngospasm. *N Z Vet J* 1975;23:148-50.
- [17] Hale GJ. Respiratory obstruction after anaesthesia. *Vet Rec* 1989;125:211-12.
- [18] Martin-Flores M, Sakai DM, Portela DA, Borlle L, Campoy L, Gleed RD. Prevention of laryngospasm with rocuronium in cats: a dose-finding study. *Vet Anaesth Analg* 2016;43:511-18.
- [19] Ead H. Post-anesthesia tracheal extubation. *Dynamics* 2004;15:20-5.
- [20] Michelet D, Skhiri A, Greff B, Luce V, Dahmani S. Management of perioperative laryngospasm by French paediatric anaesthetists. *Br J Anaesth* 2017;119:342-43.
- [21] Backus WW, Ward RR, Vitkun SA, Fitzgerald D, Askanazi J. Postextubation laryngeal spasm in an unanesthetized patient with Parkinson's disease. *J Clin Anesth* 1991;3:314-16.
- [22] Brakenhoff JE, Holcombe SJ, Hauptman JG, Smith HK, Nickels FA, Caron JP. The prevalence of laryngeal disease in a large population of competition draft horses. *Vet Surg* 2006;35:579-83.

- [23] Rooney JR, Delaney FM (1970) An hypothesis on the causation of laryngeal hemiplegia in horses. *Equine Vet J* 1970; 2:35-37.
- [24] Derksen FJ, Stick JA, Scott EA, Robinson NE, Slocumbe RF. Effect of laryngeal hemiplegia and laryngoplasty on airway flow mechanics in exercising horses. *Am J Vet Res* 1986;47:16-20.
- [25] Go L, Barton AK, Ohnesorge B. Pharyngeal diameter in various head and neck positions during exercise in sport horses. *BMC Vet Res* 2014;10:117.
- [26] Petschef VM, Derksen J, Berneyen R, Robinson E. Effect of head position on upper airway function in exercising horses. *Equine Vet J* 2010;27:18-22.
- [27] Young SS (2005) Post anaesthetic myopathy. *Equine vet. Educ* 2005;7:60-63.
- [28] Valentine BA, Cooper BJ. Incidence of Polysaccharide Storage Myopathy: Necropsy study of 225 horses. *Vet Pathol* 2005;42:823-827.
- [29] Bloom BA, Valentine BA, Gleed RD, Cable CS. Postanaesthetic recumbency in a Belgian filly with polysaccharide storage myopathy. *Vet Rec* 1999;144:73-75.
- [30] Smith RK, Dyson SJ, Head MJ, Butson RJ. Ultrasonography of the equine triceps muscle before and after general anaesthesia and in post anaesthetic myopathy. *Equine Vet J* 1996;28:311-19.
- [31] Bidwell LA, Bramlage LR, Rood WA. Equine perioperative fatalities associates with general anaesthesia at a private practice—A retrospective case series. *Vet Anaesth Analg* 2007;34:23-30.

Highlights

- Transient laryngeal dysfunction is a life-threatening peri-anesthetic complication
- Intubation and extubation may cause traumatic laryngeal dysfunction
- Draft horses might require additional care in the perioperative period

Author contributions for the manuscript entitled ‘Acute life-threatening laryngeal dysfunction in a draft horse recovering from general anesthesia: a case report’

Hayley Linda Ronaldson: Data Curation, Writing-Original Draft

Paolo Monticelli: Writing-Review & Editing, Visualization

Roger Smith: Writing-Review & Editing

Chiara Adami: Data Curation, Writing-Original Draft & Editing, Supervision

Ethical statement:

The horse object of this case report received high standard care and all the procedures carried out -including monitoring and blood sampling- are part of the routine standard care of the horse undergoing anesthesia. Data were collected from the clinical case and used retrospectively for publication.

Conflict of interest statement:

The authors declare no conflicts of interest associated with the research and preparation of this manuscript.